

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TS410245P5

Macrophage Inflammatory Protein 4 (CCL18) CytoSection

Product data:

Description:Transient overexpression of CCL18 in HEK293T cells, FFPE control for IHC, ICC and ISH staining, 25 slides per packSpecies:HumanExpression Host:HEK293TExpression cDNA Clone or AA Sequence:TrueORF Clone RC210245Tag:C-MYC/DDKDetection Antibodis:DDK Rabbit monoclonal antibody, recognizing both N- and C-terminal tags (TA592569)ACCN:MM 002988, NP 002979Synonyms:AMAC-1; AMAC1; CKb7; DCCK1; DCCK1; MIP-4; PARC; SCYA18Storage:Room TemperatureStability:HEK293T cells were transiently transfected with TrueORF cDNA plasmid, Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cultared to research use only and is not approved for use in humans or in clinical diagnosis.Refese:NP 002979Locus ID:Ga62Cytogenetics:Torq12Protein Families:Dungable Genome, Secreted ProteinOutgable Genom	Product Type:	CytoSections
Fxpression Host:HEK293TExpression cDNA ClonerueORF Clone RC210245or AA Sequence:C-MYC/DDKTag:C-MYC/DDKDetection Antibodies:DDK Rabbit monoclonal antibody, recognizing both N- and C-terminal tags (TA592569)ACCN:MM 002988, NP 002979Synonyms:AMAC-1; AMAC1; CKb7; DC-CK1; DCCK1; MIP-4; PARC; SCYA18Storage:Room TemperatureStability:Slides are guaranteed for a year from the date of receipt if proper storage instructions were followed.Preparation:HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cul- and mounted on positively charged SuperFrost slides.Note:NP 002979Locus ID:G362Cytogenetics:Trq12Protein Families:Truggable Genome, Secreted Protein	Description:	
Arb spression cDNA CloneTrueORF Clone RC210245Tag:C-MYC/DDKDetection Antibodies:DDK Rabbit monoclonal antibody, recognizing both N- and C-terminal tags (TA592569)ACCN:MM 002988, NP 002979Synonyms:AMAC-1; AMAC1; CKb7; DC-CK1; DCCK1; MIP-4; PARC; SCYA18Storage:Room TemperatureStability:Slides are guaranteed for a year from the date of receipt if proper storage instructions were followed.Preparation:HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cul- and mounted on positively charged SuperFrost slides.Note:Nb 002979Locus ID:6362Cytogenetics:17q12Protein Families:Druggable Genome, Secreted Protein	Species:	Human
or AA Sequence:Tag:C-MYC/DDKDetection Antibodies:DDK Rabbit monoclonal antibody, recognizing both N- and C-terminal tags (TA592569)ACCN:NM 002988, NP 002979Synonyms:AMAC-1; AMAC1; CKb7; DC-CK1; MIP-4; PARC; SCYA18Storage:Room TemperatureStability:Slides are guaranteed for a year from the date of receipt if proper storage instructions were followed.Preparation:HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides.Note:NP 002979Locus ID:6362Cytogenetics:17q12Protein Families:Druggable Genome, Secreted Protein	Expression Host:	HEK293T
Detection Antibodies:DDK Rabbit monoclonal antibody, recognizing both N- and C-terminal tags (TA592569)ACCN:NM 002988, NP 002979Synonyms:AMAC-1; AMAC1; CKb7; DC-CK1; DCCK1; MIP-4; PARC; SCYA18Storage:Room TemperatureStability:Slides are guaranteed for a year from the date of receipt if proper storage instructions were followed.Preparation:HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides.Note:Nb po02979Locus ID:6362Cytogenetics:17q12Protein Families:Druggable Genome, Secreted Protein	•	TrueORF Clone RC210245
ACCN:NM 002988, NP 002979Synonyms:AMAC-1; AMAC1; CKb7; DC-CK1; DCCK1; MIP-4; PARC; SCYA18Storage:Room TemperatureStability:Slides are guaranteed for a year from the date of receipt if proper storage instructions were followed.Preparation:HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides.Note:This product is for research use only and is not approved for use in humans or in clinical diagnosis.RefSeq:NP 002979Locus ID:6362Cytogenetics:17q12Protein Families:Druggable Genome, Secreted Protein	Tag:	C-MYC/DDK
Synonyms:AMAC-1; AMAC1; CKb7; DC-CK1; DCCK1; MIP-4; PARC; SCYA18Storage:Room TemperatureStability:Slides are guaranteed for a year from the date of receipt if proper storage instructions were folowed.Preparation:HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides.Note:This product is for research use only and is not approved for use in humans or in clinical diagnosis.RefSeq:NP 002979Locus ID:6362Cytogenetics:17q12Protein Families:Druggable Genome, Secreted Protein	Detection Antibodies:	DDK Rabbit monoclonal antibody, recognizing both N- and C-terminal tags (TA592569)
Storage:Room TemperatureStability:Slides are guaranteed for a year from the date of receipt if proper storage instructions were followed.Preparation:HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides.Note:This product is for research use only and is not approved for use in humans or in clinical diagnosis.RefSeq:NP 002979Locus ID:G362Cytogenetics:17q12Protein Families:Druggable Genome, Secreted Protein	ACCN:	<u>NM 002988, NP 002979</u>
Stability:Slides are guaranteed for a year from the date of receipt if proper storage instructions were followed.Preparation:HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides.Note:This product is for research use only and is not approved for use in humans or in clinical diagnosis.RefSeq:NP 002979Locus ID:6362Cytogenetics:17q12Protein Families:Druggable Genome, Secreted Protein	Synonyms:	AMAC-1; AMAC1; CKb7; DC-CK1; DCCK1; MIP-4; PARC; SCYA18
followed.Preparation:HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 μm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides.Note:This product is for research use only and is not approved for use in humans or in clinical diagnosis.RefSeq:NP 002979Locus ID:6362Cytogenetics:17q12Protein Families:Druggable Genome, Secreted Protein	Storage:	Room Temperature
were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides.Note:This product is for research use only and is not approved for use in humans or in clinical diagnosis.RefSeq:NP 002979Locus ID:6362Cytogenetics:17q12Protein Families:Druggable Genome, Secreted Protein	Stability:	
diagnosis.RefSeq:NP 002979Locus ID:6362Cytogenetics:17q12Protein Families:Druggable Genome, Secreted Protein		
Locus ID:6362Cytogenetics:17q12Protein Families:Druggable Genome, Secreted Protein	Preparation:	HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 μ m sections of the FFPE cell pellet blocks are cut
Cytogenetics:17q12Protein Families:Druggable Genome, Secreted Protein		HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 μm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides. This product is for research use only and is not approved for use in humans or in clinical
Protein Families: Druggable Genome, Secreted Protein	Note:	HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 μm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides. This product is for research use only and is not approved for use in humans or in clinical diagnosis.
	Note: RefSeq:	 HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 μm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides. This product is for research use only and is not approved for use in humans or in clinical diagnosis. <u>NP 002979</u>
Protein Pathways: Chemokine signaling pathway, Cytokine-cytokine receptor interaction	Note: RefSeq: Locus ID:	 HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 μm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides. This product is for research use only and is not approved for use in humans or in clinical diagnosis. NP 002979 6362
	Note: RefSeq: Locus ID: Cytogenetics:	 HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 μm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides. This product is for research use only and is not approved for use in humans or in clinical diagnosis. <u>NP 002979</u> 6362 17q12



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US