

Product datasheet for TS412825

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

LGR5 CytoSection

Product data:

Product Type: CytoSections

Description: Transient overexpression of LGR5 in HEK293T cells, FFPE control for IHC, ICC and ISH staining,

25 slides per pack

Species: Human
Expression Host: HEK293T

Expression cDNA Clone

or AA Sequence:

TrueORF Clone RC212825

Tag: C-MYC/DDK

Detection Antibodies: DDK Rabbit monoclonal antibody, recognizing both N- and C-terminal tags (TA592569)

Target Detection

Antibodies:

LGR5 Mouse Monoclonal Antibody [Clone ID: OTI2H7] (TA502948)

ACCN: <u>NM 003667, NP 003658</u>

Synonyms: FEX; GPR49; GPR67; GRP49; HG38

Storage: Room Temperature

Stability: Slides are guaranteed for a year from the date of receipt if proper storage instructions were

followed.

Preparation: HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells

were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin &

dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut

and mounted on positively charged SuperFrost slides.

Note: This product is for research use only and is not approved for use in humans or in clinical

diagnosis.

RefSeq: NP 003658


Locus ID: 8549

Cytogenetics: 12q21.1


Protein Families: Druggable Genome, GPCR, Transmembrane



Product images:





H&E staining of LGR5 overexpressed cytosection (TS412825), HEK293T cells were transfected with cDNA clone [RC212825], 4um sections, 10x magnification



H&E staining of LGR5 overexpressed cytosection (TS412825), HEK293T cells were transfected with cDNA clone [RC212825], 4um in thickness, 40x magnification







Immunohistology staining on LGR5 overexpressed cytosection (TS412825) with mouse anti DDK clone 11C3 ([TA180144]) at 1:1000 dilution 4C O/N. Antibody staining was achieved with HIER Tris/EDTA pH9 for 20minutes, Polink2 detection system with DAB chromogen ([D22-18]). Positive stain shown with the brown chromogen present. HEK293T cells were transfected with cDNA clone [RC212825], 4um sections, 40x magnification

Immunohistology staining on HEK293T untransfected control section screened with mouse anti-DDK clone 11C3 ([TA180144]) at 1:1000 dilution 4C O/N. Antibody staining was achieved with HIER Tris/EDTA pH9 for 20minutes, Polink2 detection system with DAB chromogen ([D22-18]). No brown chromogen showing negative stain.