

Product datasheet for **TS413132P5**


HSD17B13 CytoSection

Product data:


Product Type:	CytoSections
Description:	Transient overexpression of HSD17B13, transcript variant A, in HEK293T cells, FFPE control for IHC, ICC and ISH staining, 25 slides per pack
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	TrueORF Clone RC213132
Tag:	C-MYC/DDK
Detection Antibodies:	DDK Rabbit monoclonal antibody, recognizing both N- and C-terminal tags (TA592569)
Target Detection Antibodies:	HSD17B13 Mouse Monoclonal Antibody [Clone ID: OTI5C4] (TA814158)
ACCN:	NM_178135 , NP_835236
Synonyms:	HMFN0376; NIIL497; SCDR9; SDR16C3
Storage:	Room Temperature
Stability:	Slides are guaranteed for a year from the date of receipt if proper storage instructions were followed.
Preparation:	HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides.
Note:	This product is for research use only and is not approved for use in humans or in clinical diagnosis.
RefSeq:	NP_835236
Locus ID:	345275
Cytogenetics:	4q22.1
Protein Families:	Druggable Genome, Secreted Protein, Transmembrane




[View online »](#)

Product images:


H&E staining of HSD17B13 overexpressed cytosection ([TS413132]), HEK293T cells were transfected with cDNA clone [RC213132], 4um sections, 10x magnification



H&E staining of HSD17B13 overexpressed cytosection ([TS413132]), HEK293T cells were transfected with cDNA clone [RC213132], 4um sections, 40x magnification



Immunohistochemistry staining on HSD17B13 overexpressed cytosection ([TS413132]) with mouse anti DDK clone 11C3 ([TA180144]) at 1:1000 dilution 4C O/N. Antibody staining was achieved with HIER Tris/EDTA pH9 for 20minutes, Polink2 detection system with DAB chromogen ([D22-18]). Positive stain shown with the brown chromogen present. HEK293T cells were transfected with cDNA clone [RC213132], 4um sections, 40x magnification



Immunohistochemistry staining on HEK293T untransfected control section screened with mouse anti-DDK clone 11C3 ([TA180144]) at 1:1000 dilution 4C O/N. Antibody staining was achieved with HIER Tris/EDTA pH9 for 20minutes, Polink2 detection system with DAB chromogen ([D22-18]). No brown chromogen showing negative stain.