

## Product datasheet for **TS413631**


### CTLA4 CytoSection

#### Product data:


Product Type:	CytoSections
Description:	Transient overexpression of CTLA4, transcript variant 2, in HEK293T cells, FFPE control for IHC, ICC and ISH staining, 5 slides per pack
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	TrueORF Clone RC213631
Tag:	C-MYC/DDK
Detection Antibodies:	DDK Rabbit monoclonal antibody, recognizing both N- and C-terminal tags (TA592569)
Target Detection Antibodies:	CTLA4 Mouse Monoclonal Antibody [Clone ID: OT1G10] (TA810299)
ACCN:	<u><a href="#">NM_001037631</a></u> , <u><a href="#">NP_001032720</a></u>
Synonyms:	ALPS5; CD; CD152; CELIAC3; CTLA-4; GRD4; GSE; IDDM12
Storage:	Room Temperature
Stability:	Slides are guaranteed for a year from the date of receipt if proper storage instructions were followed.
Preparation:	HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides.
Note:	This product is for research use only and is not approved for use in humans or in clinical diagnosis.
RefSeq:	<u><a href="#">NP_001032720</a></u>
Locus ID:	1493
Cytogenetics:	2q33.2
Protein Families:	Druggable Genome, Transmembrane
Protein Pathways:	Autoimmune thyroid disease, Cell adhesion molecules (CAMs), T cell receptor signaling pathway




[View online »](#)

**Product images:**


H&E staining of CTLA4 overexpressed cytosection (TS413631), HEK293T cells were transfected with cDNA clone [RC213631], 4um sections, 10x magnification




H&E staining of CTLA4 overexpressed cytosection (TS413631), HEK293T cells were transfected with cDNA clone [RC213631], 4um in thickness, 40x magnification




Immunohistology staining on CTLA4 overexpressed cytosection (TS413631) with mouse anti CTLA4 antibody ([UM870141]) at 1:200 dilution, 1hr room temperature. The staining was achieved with HIER Tris/EDTA buffer, pH9.0, 100oC for 20 min, Polink2 detection system ([D22-18]) with DAB chromogen. HEK293T cells were transfected with cDNA clone [RC213631], 4um sections, 40x magnification



Immunohistology staining of untransfected HEK293T with mouse anti CTLA4 antibody ([UM870141]) at 1:200 dilution, 1 hour at room temperature. The staining was conducted with HIER Tris/EDTA buffer, pH9.0, 100oC for 20 min, Polink2 detection system with DAB chromogen ([D22-18]). HEK293T cells were 4um sections, 40x magnification.



Immunohistology staining on CTLA4 overexpressed cytosection (TS413631) with mouse anti DDK clone 11C3 C/N [TA180144] at 1:1000 dilution 4C O/N. Antibody staining was achieved with HIER Tris/EDTA pH9 for 20minutes, Polink2 detection system with DAB chromogen C/N [D22-18]. Positive stain shown with the brown chromogen present. HEK293T cells were transfected with cDNA clone [RC213631], 4um sections, 40x magnification



Immunohistology staining on HEK293T untransfected control section screened with mouse anti-DDK clone 11C3 C/N [TA180144] at 1:1000 dilution 4C O/N. Antibody staining was achieved with HIER Tris/EDTA pH9 for 20minutes, Polink2 detection system with DAB chromogen C/N [D22-18]. No brown chromogen showing negative stain.