

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for TS422540P5

## Metabotropic Glutamate Receptor 7 (GRM7) CytoSection

## **Product data:**

Product Type:	CytoSections
Description:	Transient overexpression of GRM7, transcript variant 2, in HEK293T cells, FFPE control for IHC, ICC and ISH staining, 25 slides per pack
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	TrueORF Clone RC222540
Tag:	C-MYC/DDK
Detection Antibodies:	DDK Rabbit monoclonal antibody, recognizing both N- and C-terminal tags (TA592569)
ACCN:	<u>NM 181874, NP 870989</u>
Synonyms:	GLUR7; GPRC1G; MGLU7; MGLUR7; NEDSHBA; PPP1R87
Storage:	Room Temperature
Stability:	Slides are guaranteed for a year from the date of receipt if proper storage instructions were
<b>,</b>	followed.
Preparation:	
-	followed. HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 μm sections of the FFPE cell pellet blocks are cut
Preparation:	followed. HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 μm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides. This product is for research use only and is not approved for use in humans or in clinical
Preparation: Note:	followed. HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 μm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides. This product is for research use only and is not approved for use in humans or in clinical diagnosis.
Preparation: Note: RefSeq:	followed. HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 μm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides. This product is for research use only and is not approved for use in humans or in clinical diagnosis. <u>NP 870989</u>
Preparation: Note: RefSeq: Locus ID:	followed. HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 μm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides. This product is for research use only and is not approved for use in humans or in clinical diagnosis. <u>NP 870989</u> 2917
Preparation: Note: RefSeq: Locus ID: Cytogenetics:	followed. HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides. This product is for research use only and is not approved for use in humans or in clinical diagnosis. <u>NP 870989</u> 2917 3p26.1



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US