

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Product datasheet for TS422881P5

## **Corticotropin Releasing Factor Receptor 2 (CRHR2) CytoSection**

## **Product data:**

Description:Transient overexpression of CRHR2, transcript variant 1, in HEK293T cells, FFPE control for HC, ICC and ISH staining, 25 slides per packSpecies:HumanExpression Host:HEK293TFxpression cDNA Clone or AA Sequence:TrueORF Clone RC222881Tag:C-MYC/DDKDetection Antibodis:DDK Rabbit monoclonal antibody, recognizing both N- and C-terminal tags (TA592569)ACCN:MM 001883, NP 001874Synonyms:CRF-RB; CRF2; CRFR2; HM-CRFStorage:Room TemperatureStability:Slides are guaranteed for a year from the date of receipt if proper storage instructions were followed.Preparation:HEK293T cells were transiently transfected with TrueORF cDNA plasmid, Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of hEFPE cell pellet blocks are cultured and mounted on positively charged SuperFrost slides.Note:NP 001874Locus ID:J395Freperict:Orugable Genome, TransmembraneProtein Families:Neuroculty ingand-receptor interaction	Product Type:	CytoSections
Fxpression Host:HEK293TExpression cDNA CloneTrueORF Clone RC222881or AA Sequence:C-MYC/DDKTag:C-MYC/DDKDetection Antibodies:DDK Rabbit monoclonal antibody, recognizing both N- and C-terminal tags (TA592569)ACCN:NM 001883, NP 001874Synonyms:CRF-RB; CRF2; CRFR2; HM-CRFStorage:Room TemperatureStability:Slides are guaranteed for a year from the date of receipt if proper storage instructions were followed.Preparation:HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cul and mounted on positively charged SuperFrost slides.Note:NP. 001874Locus ID:1395Cytogenetics:7p14.3Protein Families:Truggable Genome, Transmembrane	Description:	
Arborsion cDNA Clone or AA Sequence:TrueORF Clone RC222881Tag:C-MYC/DDKDetection Antibodis:DDK Rabbit monoclonal antibody, recognizing both N- and C-terminal tags (TA592569)ACCN:MM 001883, NP 001874Synonyms:CRF-RB; CRF2; CRFR2; HM-CRFStorage:Room TemperatureStability:Slides are guaranteed for a year from the date of receipt if proper storage instructions were followed.Preparation:HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cul- and mounted on positively charged SuperFrost slides.Note:NP 001874Gresqe:NP 001874Locus ID:1395Gytogenetics:704.3Protein Families:Druggabe Genome, Transmembrane	Species:	Human
or AA Sequence:Tag:C-MYC/DDKDetection Antibodies:DDK Rabbit monoclonal antibody, recognizing both N- and C-terminal tags (TA592569)ACCN:NM 001883, NP 001874Synonyms:CRF-RB; CRF2; CRFR2; HM-CRFStorage:Room TemperatureStability:Sildes are guaranteed for a year from the date of receipt if proper storage instructions were followed.Preparation:HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides.Note:NP 001874Locus ID:1395Cytogenetics:7p14.3Protein Families:Druggable Genome, Transmembrane	Expression Host:	HEK293T
Detection Antibodies:DDK Rabbit monoclonal antibody, recognizing both N- and C-terminal tags (TA592569)ACCN:NM 001883, NP 001874Synonyms:CRF-RB; CRF2; CRFR2; HM-CRFStorage:Room TemperatureStability:Sildes are guaranteed for a year from the date of receipt if proper storage instructions were followed.Preparation:HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cul- and mounted on positively charged SuperFrost slides.Note:Nis product is for research use only and is not approved for use in humans or in clinical diagnosis.RefSeq:NP_001874Locus ID:1395Oytogenetics:7p14.3Protein Families:Druggable Genome, Transmembrane	•	TrueORF Clone RC222881
ACCN:NM 001883, NP 001874Synonyms:CRF-RB; CRF2; CRFR2; HM-CRFStorage:Room TemperatureStability:Slides are guaranteed for a year from the date of receipt if proper storage instructions were followed.Preparation:HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides.Note:This product is for research use only and is not approved for use in humans or in clinical diagnosis.RefSeq:NP 001874Locus ID:1395Otygenetics:7p14.3Protein Families:Druggable Genome, Transmembrane	Tag:	C-MYC/DDK
Synonyms:CRF-RB; CRF2; CRFR2; HM-CRFStorage:Room TemperatureStability:Slides are guaranteed for a year from the date of receipt if proper storage instructions were folowed.Preparation:HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides.Note:This product is for research use only and is not approved for use in humans or in clinical diagnosis.RefSeq:NP 001874Locus ID:1395Otypenetics:7p14.3Protein Families:Druggabe Genome, Transmembrane	<b>Detection Antibodies:</b>	DDK Rabbit monoclonal antibody, recognizing both N- and C-terminal tags (TA592569)
Storage:Room TemperatureStability:Slides are guaranteed for a year from the date of receipt if proper storage instructions were followed.Preparation:HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides.Note:This product is for research use only and is not approved for use in humans or in clinical diagnosis.RefSeq:NP 001874Locus ID:1395Cytogenetics:7p14.3Protein Families:Druggable Genome, Transmembrane	ACCN:	<u>NM 001883</u> , <u>NP 001874</u>
Stability:Slides are guaranteed for a year from the date of receipt if proper storage instructions were followed.Preparation:HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides.Note:This product is for research use only and is not approved for use in humans or in clinical diagnosis.RefSeq:NP 001874Locus ID:1395Cytogenetics:7p14.3Protein Families:Druggable Genome, Transmembrane	Synonyms:	CRF-RB; CRF2; CRFR2; HM-CRF
followed.Preparation:HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides.Note:This product is for research use only and is not approved for use in humans or in clinical diagnosis.RefSeq:NP 001874Locus ID:1395Cytogenetics: $7$ p14.3Protein Families:Druggable Genome, Transmembrane	Storage:	Room Temperature
were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides.Note:This product is for research use only and is not approved for use in humans or in clinical diagnosis.RefSeq:NP 001874Locus ID:1395Cytogenetics:7p14.3Protein Families:Druggable Genome, Transmembrane	Stability:	
diagnosis.RefSeq:NP 001874Locus ID:1395Cytogenetics:7p14.3Protein Families:Druggable Genome, Transmembrane	Preparation:	were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 $\mu$ m sections of the FFPE cell pellet blocks are cut
Locus ID:1395Cytogenetics:7p14.3Protein Families:Druggable Genome, Transmembrane	Note:	
Cytogenetics:7p14.3Protein Families:Druggable Genome, Transmembrane	RefSeq:	<u>NP 001874</u>
Protein Families: Druggable Genome, Transmembrane	Locus ID:	1395
	Cytogenetics:	7p14.3
Protein Pathways: Neuroactive ligand-receptor interaction	Protein Families:	Druggable Genome, Transmembrane
	Protein Pathways:	Neuroactive ligand-receptor interaction



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US