

Product datasheet for **TS424141**


PD L2 (PDCD1LG2) CytoSection

Product data:


Product Type:	CytoSections
Description:	Transient overexpression of PDCD1LG2 in HEK293T cells, FFPE control for IHC, ICC and ISH staining, 25 slides per pack
Species:	Human
Expression Host:	HEK293T
Expression cDNA Clone or AA Sequence:	TrueORF Clone RC224141
Tag:	C-MYC/DDK
Detection Antibodies:	DDK Rabbit monoclonal antibody, recognizing both N- and C-terminal tags (TA592569)
Target Detection Antibodies:	PD L2 (PDCD1LG2) Mouse Monoclonal Antibody [Clone ID: OTI8G8] (TA806581)
ACCN:	NM_025239 , NP_079515
Synonyms:	B7DC; bA574F11.2; Btdc; CD273; PD-L2; PDCD1L2; PDL2
Storage:	Room Temperature
Stability:	Slides are guaranteed for a year from the date of receipt if proper storage instructions were followed.
Preparation:	HEK293T cells were transiently transfected with TrueORF cDNA plasmid. Transfected cells were cultured for 48hrs. After harvesting, the cultured cells were fixed in formalin & dehydrated before embedding in paraffin. 5 µm sections of the FFPE cell pellet blocks are cut and mounted on positively charged SuperFrost slides.
Note:	This product is for research use only and is not approved for use in humans or in clinical diagnosis.
RefSeq:	NP_079515
Locus ID:	80380
Cytogenetics:	9p24.1
Protein Families:	Transmembrane
Protein Pathways:	Cell adhesion molecules (CAMs)




[View online »](#)

Product images:


H&E staining of PD-L2 overexpressed cytosection (TS424141), HEK293T cells were transfected with cDNA clone [RC202141], 4um in thickness, 40x magnification




Immunohistochemistry staining on PD-L2 overexpressed cytosection (TS424141) with mouse anti PD-L2 antibody ([UM800115]) at 1:200 dilution, 1hr room temperature. The staining was achieved with HIER Tris/EDTA buffer, pH9.0, 100oC for 20 min, Polink2 detection system ([D22-18]) with DAB chromogen. HEK293T cells were transfected with cDNA clone [RC202141], 4um sections, 40x magnification



Immunohistology staining of untransfected HEK293T with mouse anti PD-L2 antibody ([UM800115]) at 1:200 dilution, 1 hour at room temperature. The staining was conducted with HIER Tris/EDTA buffer, pH9.0, 100oC for 20 min, Polink2 detection system with DAB chromogen ([D22-18]). HEK293T cells were 4um sections, 40x magnification.



Immunohistology staining on PD-L2 overexpressed cytosection (TS424141) with mouse anti DDK clone 11C3 C/N [TA180144] at 1:1000 dilution 4C O/N. Antibody staining was achieved with HIER Tris/EDTA pH9 for 20minutes, Polink2 detection system with DAB chromogen ([D22-18]). Positive stain shown with the brown chromogen present. HEK293T cells were transfected with cDNA clone [RC202141], 4um sections, 40x magnification



Immunohistology staining on HEK293T untransfected control section screened with mouse anti-DDK clone 11C3 ([TA180144]) at 1:1000 dilution 4C O/N. Antibody staining was achieved with HIER Tris/EDTA pH9 for 20minutes, Polink2 detection system with DAB chromogen ([D22-18]). No brown chromogen showing negative stain.