

Product datasheet for **UM50007**

BubR1 (BUB1B) Mouse Monoclonal Antibody [Clone ID: UMAB7]

Product data:

Product Type:	Primary Antibodies
Clone Name:	UMAB7
Applications:	IF, IHC, WB
Recommended Dilution:	WB 1:2000, IHC 1:50, IF 1:100, Flow 1:100, IF 1:100
Reactivity:	Human, Monkey
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Protein expressed in 293T cell transfected with human BUB1B expression vector
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.5~1.0 mg/ml (Lot Dependent)
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	119.5 kDa
Gene Name:	BUB1 mitotic checkpoint serine/threonine kinase B
Database Link:	NP_001202 Entrez Gene 100423079 Monkey Entrez Gene 701 Human O60566
Background:	This gene encodes a kinase involved in spindle checkpoint function. The protein has been localized to the kinetochore and plays a role in the inhibition of the anaphase-promoting complex/cyclosome(APC/C), delaying the onset of anaphase and ensuring proper chromosome segregation. Impaired spindle checkpoint function has been found in many forms of cancer.
Synonyms:	Bub1A; BUB1beta; BUBR1; hBUBR1; MAD3L; MVA1; SSK1




[View online »](#)


Protein Families: Druggable Genome, Protein Kinase, Stem cell - Pluripotency

Protein Pathways: Cell cycle


Product images:




Immunohistochemical staining of paraffin-embedded Carcinoma of lung tissue using anti-BUB1B mouse monoclonal antibody. (Clone UMAB7, dilution 1:100; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)




Immunohistochemical staining of paraffin-embedded human endometrial carcinoma using anti-BUB1B a mouse monoclonal antibody. Heat-induced epitope retrieval by ACCEL (pH8.7) in pressure chamber/cooker at 110°C for 3min, UM500007 clone UMAB7 at 1:250. Images shows tumor cells with strong cytoplasmic staining.




Immunohistochemical staining of paraffin-embedded mouse liver tissue using anti-BUB1B clone UMAB7 mouse monoclonal antibody. HIER TEE buffer pH9 ([B21-100]) at 110C for 10 min, UM500007 (1:100). Detection was done with Klear Mouse (C/N [D52-18]) DAB Kit.




Immunohistochemical staining of paraffin-embedded mouse lung tissue using anti-BUB1B clone UMAB7 mouse monoclonal antibody. HIER TEE buffer pH9 ([B21-100]) at 110C for 10 min, UM500007 (1:100). Detection was done with Klear Mouse (C/N [D52-18]) DAB Kit.




Immunohistochemical staining of paraffin-embedded mouse spleen tissue using anti-BUB1B clone UMAB7 mouse monoclonal antibody. HIER TEE buffer pH9 ([B21-100]) at 110C for 10 min, UM500007 (1:100). Detection was done with Klear Mouse (C/N [D52-18]) DAB Kit.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-BUB1B monoclonal antibody (Clone UMAB7) at 1:500.



Immunofluorescent staining of A549 cells using anti-BUB1B mouse monoclonal antibody (UM500007, green). Actin filaments were labeled with TRITC-phalloidin (red), and nuclear with DAPI (blue). The three-color overlay image is located at the bottom-right corner.



Immunofluorescent staining of HeLa cells using anti-BUB1B mouse monoclonal antibody (UM500007, green). Actin filaments were labeled with TRITC-phalloidin (red), and nuclear with DAPI (blue). The three-color overlay image is located at the bottom-right corner.