

Product datasheet for **UM500031**

CD31 (PECAM1) Mouse Monoclonal Antibody [Clone ID: UMAB31]


Product data:

Product Type:	Primary Antibodies
Clone Name:	UMAB31
Applications:	FC, IHC, WB
Recommended Dilution:	WB 1:500, IHC 1:100
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human PECAM1(NP_000433) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.5~1.0 mg/ml (Lot Dependent)
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	82.4 kDa
Gene Name:	platelet and endothelial cell adhesion molecule 1
Database Link:	NP_000433 Entrez Gene 5175 Human P16284
Background:	The protein encoded by this gene is found on the surface of platelets, monocytes, neutrophils, and some types of T-cells, and makes up a large portion of endothelial cell intercellular junctions. The encoded protein is a member of the immunoglobulin superfamily and is likely involved in leukocyte migration, angiogenesis, and integrin activation. [provided by RefSeq]




[View online »](#)


Synonyms: CD31; CD31/EndoCAM; endoCAM; GPIIA'; PECA1; PECAM-1
Protein Families: Druggable Genome, ES Cell Differentiation/IPS, Transmembrane
Protein Pathways: Cell adhesion molecules (CAMs), Leukocyte transendothelial migration

Product images:


Immunohistochemical staining of paraffin-embedded Adenocarcinoma of colon tissue using anti-PECAM1 mouse monoclonal antibody. (Clone UMAB31, dilution 1:100; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)




Immunohistochemical staining of paraffin-embedded Carcinoma of liver tissue using anti-PECAM1 mouse monoclonal antibody. (Clone UMAB31, dilution 1:100; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)




Immunohistochemical staining of paraffin-embedded pancreas tissue using anti-PECAM1 mouse monoclonal antibody. (Clone UMAB31, dilution 1:100; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)




Immunohistochemical staining of paraffin-embedded Carcinoma of thyroid tissue using anti-PECAM1 mouse monoclonal antibody. (Clone UMAB31, dilution 1:100; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)




Immunohistochemical staining of paraffin-embedded endometrium tissue using anti-PECAM1 mouse monoclonal antibody. (Clone UMAB31, dilution 1:100; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of endometrium tissue using anti-PECAM1 mouse monoclonal antibody. (Clone UMAB31, dilution 1:100; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)




Immunohistochemical staining of paraffin-embedded prostate tissue using anti-PECAM1 mouse monoclonal antibody. (Clone UMAB31, dilution 1:100; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)




Immunohistochemical staining of paraffin-embedded Carcinoma of prostate tissue using anti-PECAM1 mouse monoclonal antibody. (Clone UMAB31, dilution 1:100; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)




Immunohistochemical staining of paraffin-embedded Carcinoma of bladder tissue using anti-PECAM1 mouse monoclonal antibody. (Clone UMAB31, dilution 1:100; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)




Immunohistochemical staining of paraffin-embedded human placenta using anti-PECAM1 mouse monoclonal antibody (UM500031) at 1:100 with Polink2 Broad HRP DAB detection kit; heat-induced epitope retrieval with GBI Citrate pH6.0 HIER buffer using pressure chamber for 3 minutes at 110C. Cytoplasmic and membranous staining is seen in the endothelia cells and weak staining in the trophoblast.




Immunohistochemical staining of paraffin-embedded human skin using anti-PECAM1 mouse monoclonal antibody (UM500031) at 1:100 with Polink2 Broad HRP DAB detection kit; heat-induced epitope retrieval with GBI Citrate pH6.0 HIER buffer using pressure chamber for 3 minutes at 110C. Cytoplasmic and membranous staining is seen in the vasular endothelia cells.




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PECAM1 ([RC208654], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PECAM1 mouse monoclonal antibody.1:500`



Western blot analysis of extracts (35ug) from 10 different human tissues and Jurkat cell lines by using anti-PECAM1 monoclonal antibody. 1:250



Western Blot analysis of 10 different human tissue lysates (10ug) by using anti-PECAM1 monoclonal antibody (clone UMAB31, 1:500)



Living HEK293T cells transfected with either [RC208654] plasmid (red) or empty vector (blue) were immunostained by anti-PECAM1 antibody (UM500031) or isotype control antibody (green), and then analyzed by flow cytometry (1:100).