

Product datasheet for **UM500049CF**

p53 (TP53) Mouse Monoclonal Antibody [Clone ID: UMAB55]

Product data:

| | |
|-------------------------|--|
| Product Type: | Primary Antibodies |
| Clone Name: | UMAB55 |
| Applications: | IF, IHC, WB |
| Recommended Dilution: | IHC 1:100 |
| Reactivity: | Human |
| Host: | Mouse |
| Isotype: | IgG1 |
| Clonality: | Monoclonal |
| Immunogen: | Full length human recombinant protein of human TP53 (NP_000537) produced in HEK293T cell. |
| Formulation: | Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose) |
| Reconstitution Method: | For reconstitution, we recommend adding 100uL distilled water to a final antibody concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific) |
| Purification: | Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G) |
| Conjugation: | Unconjugated |
| Storage: | Store at -20°C as received. |
| Stability: | Stable for 12 months from date of receipt. |
| Predicted Protein Size: | 43.5 kDa |
| Gene Name: | tumor protein p53 |
| Database Link: | NP_000537 Entrez Gene 7157 Human P04637 |



[View online »](#)

Background:

The protein encoded by this gene is a member of the STAT family of transcription factors. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is activated by, and mediates the responses of many cell ligands, such as IL2, IL3, IL7 GM-CSF, erythropoietin, thrombopoietin, and different growth hormones. Activation of this protein in myeloma and lymphoma associated with a TEL/JAK2 gene fusion is independent of cell stimulus and has been shown to be essential for the tumorigenesis. The mouse counterpart of this gene is found to induce the expression of BCL2L1/BCL-X(L), which suggests the antiapoptotic function of this gene in cells. [provided by RefSeq, Jul 2008]

Synonyms:

BCC7; LFS1; P53; TRP53

Protein Families:


Druggable Genome, Stem cell - Pluripotency, Transcription Factors

Protein Pathways:


Amyotrophic lateral sclerosis (ALS), Apoptosis, Basal cell carcinoma, Bladder cancer, Cell cycle, Chronic myeloid leukemia, Colorectal cancer, Endometrial cancer, Glioma, Huntington's disease, MAPK signaling pathway, Melanoma, Neurotrophin signaling pathway, Non-small cell lung cancer, p53 signaling pathway, Pancreatic cancer, Pathways in cancer, Prostate cancer, Small cell lung cancer, Thyroid cancer, Wnt signaling pathway

Product images:

Normal Colon




Colorectal Cancer




UM500049

Immunohistochemical staining of paraffin-embedded Human normal colon tissue and colonrectal cancer tissue using anti-TP53 mouse monoclonal antibody. ([UM500049]; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)

Normal lung




Lung Adenocarcinoma




UM500049

Immunohistochemical staining of paraffin-embedded Human normal lung tissue and lung adenocarcinoma tissue using anti-TP53 mouse monoclonal antibody. ([UM500049]; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)

Normal Ovary




Ovarian Cancer




UM500049

Immunohistochemical staining of paraffin-embedded Human normal ovary tissue and ovary cancer tissue using anti-TP53 mouse monoclonal antibody. ([UM500049]; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)

Normal Breast




Breast Adenocarcinoma




UM500049


Immunohistochemical staining of paraffin-embedded Human normal breast tissue and breast adenocarcinoma tissue using anti-TP53 mouse monoclonal antibody. ([UM500049]; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)




Immunohistochemical staining of paraffin-embedded mouse testis tissue using anti-p53 clone UMAB55 mouse monoclonal antibody. HIER TEE buffer pH9 ([B21-100]) at 110C for 10 min, [UM500049] (1:100). Detection was done with Klear Mouse (C/N [D52-18]) DAB Kit.



Western Blot analysis of extracts (35ug) from 11 different cell lines by using anti-TP53 monoclonal antibody (Clone UMAB55)



Immunofluorescent staining of COS7 cells using TP53 mouse monoclonal antibody ([UM500049], green). Actin filaments were labeled with TRITC-phalloidin (red), and nuclear with DAPI (blue). The three-color overlay image is located at the bottom-right corner.



Immunofluorescent staining of HT-29 cells using TP53 mouse monoclonal antibody ([UM500049], green). Actin filaments were labeled with TRITC-phalloidin (red), and nuclear with DAPI (blue). The three-color overlay image is located at the bottom-right corner.