

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for UM500053

p53 (TP53) Mouse Monoclonal Antibody [Clone ID: UMAB62]

Product data:


Product Type:	Primary Antibodies
Clone Name:	UMAB62
Applications:	10k-ChIP, IF, IHC, WB
Recommended Dilution:	IHC 1:50~100
Reactivity:	Human
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human TP53 (NP_000537) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.5~1.0 mg/ml (Lot Dependent)
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	43.5 kDa
Gene Name:	tumor protein p53
Database Link:	<u>NP 000537</u>
	<u>Entrez Gene 7157 Human</u> <u>P04637</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	p53 (TP53) Mouse Monoclonal Antibody [Clone ID: UMAB62] – UM500053
Background:	The protein encoded by this gene is a member of the STAT family of transcription factors. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is activated by, and mediates the responses of many cell ligands, such as IL2, IL3, IL7 GM-CSF, erythropoietin, thrombopoietin, and different growth hormones. Activation of this protein in myeloma and lymphoma associated with a TEL/JAK2 gene fusion is independent of cell stimulus and has been shown to be essential for the tumorigenesis. The mouse counterpart of this gene is found to induce the expression of BCL2L1/BCL-X(L), which suggests the antiapoptotic function of this gene in cells. [provided by RefSeq, Jul 2008]
Synonyms:	BCC7; LFS1; P53; TRP53
Protein Families:	Druggable Genome, Stem cell - Pluripotency, Transcription Factors
Protein Pathway	Amyotrophic lateral sclerosis (ALS), Apoptosis, Basal cell carcinoma, Bladder cancer, Cell cycle, Chronic myeloid leukemia, Colorectal cancer, Endometrial cancer, Glioma, Huntington's disease, MAPK signaling pathway, Melanoma, Neurotrophin signaling pathway, Non-small cell lung cancer, p53 signaling pathway, Pancreatic cancer, Pathways in cancer, Prostate cancer, Small cell lung cancer, Thyroid cancer, Wnt signaling pathway


Product images:



Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human breast tissue using anti-TP53 mouse monoclonal antibody. (UM500053; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US 


Immunohistochemical staining of paraffinembedded Human normal colon tissue and colorectal cancer tissue using anti-TP53 mouse monoclonal antibody. (UM500053; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)



embedded Human normal endometrium tissue and endometrium adenocarcinoma tissue using anti-TP53 mouse monoclonal antibody. (UM500053; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)


Immunohistochemical staining of paraffin-

HepG2 HeLa SV-T2 A549 CO S7 Jurkat MDCK PC-12 MCF7 BT549 HT-29




Western Blot analysis of extracts (35ug) from 11 different cell lines by using anti-TP53 monoclonal antibody (Clone UMAB62)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Immunofluorescent staining of COS7 cells using TP53 mouse monoclonal antibody (UM500053, green). Actin filaments were labeled with TRITCphalloidin (red), and nuclear with DAPI (blue). The three-color overlay image is located at the bottom-right corner.

Immunofluorescent staining of HT-29 cells using TP53 mouse monoclonal antibody (UM500053, green). Actin filaments were labeled with TRITCphalloidin (red), and nuclear with DAPI (blue). The three-color overlay image is located at the bottom-right corner.



OriGene overexpression protein microarray chip was immunostained with UltraMAB anti-TP53 mouse monoclonal antibody (UM500053). The positive reactive proteins are highlighted with two red arrows in the enlarged subarray. All the positive controls spotted in this subarray are also labeled for clarification.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US