

Product datasheet for UM500068

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Kallikrein 8 (KLK8) Mouse Monoclonal Antibody [Clone ID: UMAB84]

Product data:

Product Type: Primary Antibodies

Clone Name: UMAB84
Applications: IHC, WB
Recommended Dilution: IHC 1:100
Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human KLK8(NP_009127) produced in HEK293T

cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5~1.0 mg/ml (Lot Dependent)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 27.9 kDa

Gene Name: kallikrein related peptidase 8

Database Link: NP 009127

Entrez Gene 11202 Human

060259






Background:

Kallikreins are a subgroup of serine proteases having diverse physiological functions. Growing evidence suggests that many kallikreins are implicated in carcinogenesis and some have potential as novel cancer and other disease biomarkers. This gene is one of the fifteen kallikrein subfamily members located in a cluster on chromosome 19. Alternate splicing of this gene results in four transcript variants encoding four different isoforms. The isoforms exhibit distinct patterns of expression that suggest roles in brain plasticity and ovarian cancer. [provided by RefSeq, Jul


Synonyms: HNP; NP; NRPN; PRSS19; TADG14

Protein Families: Druggable Genome, Secreted Protein, Transmembrane

Product images:




Immunohistochemical staining of paraffinembedded Human embryonic brain cortex tissue using anti-KLK8 mouse monoclonal antibody. (UM500068; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)




Western blot analysis of extracts (10ug) from 10 Human tissue by using anti-KLK8 monoclonal antibody at 1:500 (1: Testis; 2: Omentum; 3: Uterus; 4: Breast; 5: Brain; 6: Liver; 7: Ovary; 8: Thyroid gland; 9: colon;10: spleen).





Western blot of cell lysates (35ug) from 9 different cell lines (1: HepG2, 2: HeLa, 3: SV-T2, 4: A549, 5: COS7, 6: Jurkat, 7: MDCK, 8: PC-12, 9: MCF7). Diluation: 1:500



Western blot of mouse tissue lysates (20ug) from Uterus. Primary antibody diluation: 1:500. Secondary antibody dilution: Mouse TrueBlot® Ultra (1:1000).