

Product datasheet for UM570021

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

XPF (ERCC4) Mouse Monoclonal Antibody [Clone ID: UMAB21]

Product data:

Product Type: Primary Antibodies

Clone Name: UMAB21

Applications: 10k-ChIP, IF, IHC, WB

Recommended Dilution: WB 1:200~500, IF 1:100, FLOW 1:100, IHC 1:500

Reactivity: Human, Dog, Rat, Monkey, Mouse

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human XPF (NP_005227) produced in HEK293 cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5~1.0 mg/ml (Lot Dependent)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 104.3 kDa

Gene Name: ERCC excision repair 4, endonuclease catalytic subunit

Database Link: NP 005227

Entrez Gene 50505 MouseEntrez Gene 304719 RatEntrez Gene 479842 DogEntrez Gene

712641 MonkeyEntrez Gene 2072 Human

O92889

Background: The protein encoded by this gene forms a complex with ERCC1 and is involved in the 5'

incision made during nucleotide excision repair. This complex is a structure specific DNA repair endonuclease that interacts with EME1. Defects in this gene are a cause of xeroderma

pigmentosum complementation group F (XP-F), or xeroderma pigmentosum VI (XP6).

Synonyms: ERCC11; FANCQ; RAD1; XFEPS; XPF







Protein Families: Druggable Genome

Protein Pathways: Nucleotide excision repair


Product images:



Immunohistochemical staining of paraffinembedded Carcinoma of lung tissue using anti-XPF (UMAB21) mouse monoclonal antibody. ([UM500021], Dilution 1:50; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)




Western blot analysis of crude cell extracts (35ug) from 9 different cell lines by using anti-XPF (UMAB21) monoclonal antibody.




Western Blot analysis of 10 different human tissue lysates (10ug) by using anti-XPF monoclonal antibody (clone UMAB21, 1:500)





Anti-XPF (UMAB21) mouse monoclonal antibody ([UM500021]) immunofluorescent staining of COS7 cells transiently transfected by pCMV6-ENTRY XPF ([RC223300]).



OriGene overexpression protein microarray chip was immunostained with UltraMAB anti-XPF mouse monoclonal antibody (clone UMAB21). The positive reactive proteins are highlighted with two red arrows in the enlarged subarray. All the positive controls spotted in this subarray are also labeled for clarification. These data show that UltraMAB anti-XPF (UMAB21) very specifically recognizes XPF antigen on OriGene protein microarray chip.