

Product datasheet for UM570081

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ZNF447 (ZSCAN18) Mouse Monoclonal Antibody [Clone ID: UMAB114]

Product data:

Product Type: Primary Antibodies

Clone Name: UMAB114

Applications: 10k-ChIP, IHC, WB

Recommended Dilution: IHC 1:100~200

Reactivity: Human Host: Mouse

Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human ZSCAN18(NP_076415) produced in

HEK293T cell.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5~1.0 mg/ml (Lot Dependent)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 54.6 kDa

Gene Name: zinc finger and SCAN domain containing 18

Database Link: NP 076415

Entrez Gene 65982 Human

Q8TBC5


Synonyms: ZNF447

Protein Families: Transcription Factors






Product images:




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human breast tissue using anti-ZSCAN18 mouse monoclonal antibody. ([UM500081]; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)




Immunohistochemical staining of paraffinembedded Human Kidney tissue using anti-ZSCAN18 mouse monoclonal antibody. ([UM500081]; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)






Immunohistochemical staining of paraffinembedded Carcinoma of Human kidney tissue using anti-ZSCAN18 mouse monoclonal antibody. ([UM500081]; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)




Immunohistochemical staining of paraffinembedded Carcinoma of Human liver tissue using anti-ZSCAN18 mouse monoclonal antibody. ([UM500081]; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)




Immunohistochemical staining of paraffinembedded Carcinoma of Human lung tissue using anti-ZSCAN18mouse monoclonal antibody. ([UM500081]; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)






Immunohistochemical staining of paraffinembedded Human pancreas tissue using anti-ZSCAN18 mouse monoclonal antibody. ([UM500081]; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)




Immunohistochemical staining of paraffinembedded Carcinoma of Human pancreas tissue using anti-ZSCAN18 mouse monoclonal antibody. ([UM500081]; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)




Immunohistochemical staining of paraffinembedded Human thyroid tissue using anti-ZSCAN18 mouse monoclonal antibody. ([UM500081]; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)






Immunohistochemical staining of paraffinembedded Carcinoma of Human thyroid tissue using anti-ZSCAN18 mouse monoclonal antibody. ([UM500081]; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)




Immunohistochemical staining of paraffinembedded Human endometrium tissue using anti-ZSCAN18 mouse monoclonal antibody. ([UM500081]; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)




Immunohistochemical staining of paraffinembedded Carcinoma of Human bladder tissue using anti-ZSCAN18 mouse monoclonal antibody. ([UM500081]; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)






Immunohistochemical staining of paraffinembedded Human lymph node tissue using anti-ZSCAN18 mouse monoclonal antibody. ([UM500081]; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)




Western blot of cell lysates (35ug) from 9 different cell lines (1: HepG2, 2: HeLa, 3: SV-T2, 4: A549. 5: COS7, 6: Jurkat, 7: MDCK, 8: PC-12, 9: MCF7).




Western blot of human tissue lysates (15ug) from 10 different tissues (1: Testis, 2: Omentum, 3: Uterus, 4: Breast, 5: Brain, 6: Liver, 7: Ovary, 8: Colon, 9: Spleen, 10: Thyroid). Diluation: 1:500.






Immunohistochemical staining of paraffinembedded mouse kidney using mouse anti-ZSCAN18 clone UMAB114 mouse monoclonal antibody. HIER TEE buffer ([B21-100]) (pH9) at 110C for 10 min, [UM500081] (1:100). Detection was done with Klear Mouse (C/N [D52-18]) DAB Chromogen Kit.




Immunohistochemical staining of paraffinembedded mouse brain using mouse anti-ZSCAN18 clone UMAB114 mouse monoclonal antibody. HIER TEE buffer ([B21-100]) (pH9) at 110C for 10 min, [UM500081] (1:100). Detection was done with Klear Mouse (C/N [D52-18]) DAB Chromogen Kit.



Immunohistochemical staining of paraffinembedded mouse lung using mouse anti-ZSCAN18 clone UMAB114 mouse monoclonal antibody. HIER TEE buffer ([B21-100]) (pH9) at 110C for 10 min, [UM500081] (1:100). Detection was done with Klear Mouse (C/N [D52-18]) DAB Chromogen Kit.





OriGene overexpression protein microarray chip was immunostained with UltraMAB anti-ZSCAN18 mouse monoclonal antibody ([UM500081]). The positive reactive proteins are highlighted with two red arrows in the enlarged subarray. All the positive controls spotted in this subarray are also labeled for clarification.