

## **Product datasheet for UM570083**

# OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# **ALX4 Mouse Monoclonal Antibody [Clone ID: UMAB118]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: UMAB118

**Applications:** 10k-ChIP, IHC, WB **Recommended Dilution:** IHC 1:100~200

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human ALX4(NP\_068745) produced in HEK293T

cell.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.5~1.0 mg/ml (Lot Dependent)

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 44.1 kDa

Gene Name: ALX homeobox 4

Database Link: NP 068745

Entrez Gene 11695 MouseEntrez Gene 296511 RatEntrez Gene 60529 Human

O9H161






Background:

This gene encodes a paired-like homeodomain transcription factor expressed in the mesenchyme of developing bones, limbs, hair, teeth, and mammary tissue. Mutations in this gene cause parietal foramina 2 (PFM2); an autosomal dominant disease characterized by deficient ossification of the parietal bones. Mutations in this gene also cause a form of frontonasal dysplasia with alopecia and hypogonadism; suggesting a role for this gene in craniofacial development, mesenchymal-epithelial communication, and hair follicle development. Deletion of a segment of chromosome 11 containing this gene, del(11)(p11p12), causes Potocki-Shaffer syndrome (PSS); a syndrome characterized by craniofacial anomalies, mental retardation, multiple exostoses, and genital abnormalities in males. In mouse, this gene has been shown to use dual translation initiation sites located 16 codons apart. [provided by RefSeq, Oct 2009]

Synonyms: CRS5; FND2


**Protein Families:** Druggable Genome

# **Product images:**




Immunohistochemical staining of FFPE human breast tissue section using anti-ALX4 mouse monoclonal antibody [UM500083], clone UMAB118 at 1:200. (Heat-induced epitope retrieval by Tris-EDTA, pH9.0; Polink1 Broad HRP for 15 min; DAB chromogen for 5 min & DAB enhancer for 30 sec).





OriGene overexpression protein microarray chip was immunostained with UltraMAB anti-ALX4 mouse monoclonal antibody ([UM500083]). The positive reactive proteins are highlighted with two red arrows in the enlarged subarray. All the positive controls spotted in this subarray are also labeled for clarification.



Western blot of cell lysates (35ug) from 8 different cell lines (1: HepG2, 2: HeLa, 3: SV-T2, 4: MCF7, 5: COS7, 6: Jurkat, 7: MDCK, 8: PC-12). Diluation: 1:500.