

## Product datasheet for **UM570092**

### MSI1 Mouse Monoclonal Antibody [Clone ID: UMAB127]

#### Product data:


Product Type:	Primary Antibodies
Clone Name:	UMAB127
Applications:	10k-ChIP, IHC
Recommended Dilution:	IHC 1:100~200
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human MSI1(NP_002433) produced in HEk293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.5~1.0 mg/ml (Lot Dependent)
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	38.9 kDa
Gene Name:	Homo sapiens musashi RNA binding protein 1 (MSI1), mRNA.
Database Link:	<a href="#">NP_002433</a> <a href="#">Entrez Gene 17690 Mouse</a> <a href="#">Entrez Gene 259272 Rat</a> <a href="#">Entrez Gene 4440 Human</a> <a href="#">O43347</a>
Background:	This gene encodes a protein containing two conserved tandem RNA recognition motifs. Similar proteins in other species function as RNA-binding proteins and play central roles in posttranscriptional gene regulation. Expression of this gene has been correlated with the grade of the malignancy and proliferative activity in gliomas and melanomas. A pseudogene for this gene is located on chromosome 11q13. [provided by RefSeq, Jul 2008]




[View online »](#)

Synonyms: musashi 1; musashi homolog 1 (Drosophila)


**Product images:**




Immunohistochemical staining of paraffin-embedded Human adult brain tissue using anti-MSI1 mouse monoclonal antibody. ([UM500092]; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)




Immunohistochemical staining of paraffin-embedded Human embryonic brain cortex tissue using anti-MSI1 mouse monoclonal antibody. ([UM500092]; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)




Immunohistochemical staining of paraffin-embedded Human embryonic cerebellum using anti-MS11 mouse monoclonal antibody. [UM500092]; heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min)



Immunohistochemical staining of paraffin-embedded human colon using anti-MS11 clone UMAB127 mouse monoclonal antibody at 1:200 dilution 1mg/mL and detection with Polink2 Broad HRP DAB. [UM500092] requires heat-induced epitope retrieval with Accel pH 8.7 in a pressure cooker for 3 minutes at 110C.



Immunohistochemical staining of paraffin-embedded human pancreas using anti-MS11 clone UMAB127 mouse monoclonal antibody at 1:200 dilution 1mg/mL and detection with Polink2 Broad HRP DAB. [UM500092] requires heat-induced epitope retrieval with Accel pH 8.7 in a pressure cooker for 3 minutes at 110C. Cytoplasmic and membrane staining seen weakly on the glandular cells and strongly on the Islet of Langerhans



OriGene overexpression protein microarray chip was immunostained with UltraMAB anti-MSI1 mouse monoclonal antibody ([UM500092]). The positive reactive proteins are highlighted with two red arrows in the enlarged subarray. All the positive controls spotted in this subarray are also labeled for clarification.