

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for UM800073CF

C14orf166 (RTRAF) Mouse Monoclonal Antibody [Clone ID: UMAB181]


Product data:

Product Type:	Primary Antibodies
Clone Name:	UMAB181
Applications:	10k-ChIP, IF, IHC, WB
Recommended Dilution:	IHC 1:100~200
Reactivity:	Human
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human C14orf166 (NP_057123) produced in E.coli.
Formulation:	Lyophilized powder (original buffer 1X PBS, pH 7.3, 8% trehalose)
Reconstitution Method:	For reconstitution, we recommend adding 100uL distilled water to a final antibody concentration of about 1 mg/mL. To use this carrier-free antibody for conjugation experiment, we strongly recommend performing another round of desalting process. (OriGene recommends Zeba Spin Desalting Columns, 7KMWCO from Thermo Scientific)
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	27.9 kDa
Gene Name:	RNA transcription, translation and transport factor
Database Link:	<u>NP_057123</u> <u>Entrez Gene 51637 Human</u> <u>Q9Y224</u>
Synonyms:	CGI-99; CGI99; CLE; CLE7; hCLE1; LCRP369; RLLM1




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US


Product images:



Immunohistochemical staining of paraffinembedded Carcinoma of Human prostate tissue using anti-C14orf166 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.0) at 110°C for 10min, [UM800073]) (1:200)



Immunohistochemical staining of paraffinembedded Human adult heart tissue using anti-C14orf166 mouse monoclonal antibody. (Heatinduced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.0) at 110°C for 10min, [UM800073]) (1:200)

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US 


Immunohistochemical staining of paraffinembedded human colon cancer using anti-C14ORF166 clone UMAB181 mouse monoclonal antibody at 1:600 dilution of 1mg/mL and detection with Polink2 Broad HRP DAB. [UM800073] requires heat-induced epitope retrieval with Accel for 3minutes at110C in pressure chamber. The image shows the tumor cells have strong nuclear and weak cytoplasmic staining.

Immunohistochemical staining of paraffinembedded human thyroid using anti-C14ORF166 clone UMAB181 mouse monoclonal antibody at 1:600 dilution of 1mg/mL and detection with Polink2 Broad HRP DAB. [UM800073] requires heat-induced epitope retrieval with Accel for 3minutes at110C in pressure chamber. The image shows strong nuclear and intermediate levels of cytoplasmic staining on the thyroid.


Immunohistochemical staining of paraffinembedded human kidney using anti-C14ORF166 clone UMAB181 mouse monoclonal antibody at 1:600 dilution of 1mg/mL and detection with Polink2 Broad HRP DAB. [UM800073] requires heat-induced epitope retrieval with Accel for 3minutes at110C in pressure chamber. The image shows the tumor cells have strong nuclear and intermediate levels of cytoplasmic staining.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US






Western blot of cell lysates (35ug) from 9 different cell lines (1: HepG2, 2: HeLa, 3: SV-T2, 4: A549. 5: Jurkat, 6: MDCK, 7: COS7, 8: PC-12, 9: MCF7).




Western blot analysis of extracts (15ug) from 9 Human tissue by using anti-C14orf166 monoclonal antibody (1: Testis; 2: Uterus; 3: Breast; 4: Brain; 5: Liver; 6: Ovary; 7: Thyroid gland; 8: colon;;9:Spleen). (1:500) Dilution: 1:500



Immunofluorescent staining of HeLa cells using anti-C14orf166 mouse monoclonal antibody ([UM800073], green, 1:100). Actin filaments were labeled with Alexa Fluor® 594 Phalloidin (red), and nuclear with DAPI (blue).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



OriGene overexpression protein microarray chip was immunostained with UltraMAB anti-C14orf166 mouse monoclonal antibody ([UM800073]). The positive reactive proteins are highlighted with two red arrows in the enlarged subarray. All the positive controls spotted in this subarray are also labeled for clarification (1:100).

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2024 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US