

Product datasheet for **UM800124**


DHLAG (CD74) Mouse Monoclonal Antibody [Clone ID: UMAB232]

Product data:


Product Type:	Primary Antibodies
Clone Name:	UMAB232
Applications:	IHC, WB
Recommended Dilution:	WB 1:500, IHC 1:800
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human CD74(NP_001020330) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.5~1.0 mg/ml (Lot Dependent)
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	33.3 kDa
Gene Name:	CD74 molecule
Database Link:	NP_001020330 Entrez Gene 972 Human P04233
Synonyms:	DHLAG; HLADG; Ia-GAMMA; II
Protein Families:	Druggable Genome, Transmembrane
Protein Pathways:	Antigen processing and presentation




[View online »](#)

Product images:


Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human colon tissue using anti-CD74 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.0) at 120°C for 2.5 min, UM800124) (1:800)




Immunohistochemical staining of paraffin-embedded Carcinoma of Human liver tissue using anti-CD74 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.0) at 120°C for 2.5 min, UM800124) (1:800)




Immunohistochemical staining of paraffin-embedded Carcinoma of Human lung tissue using anti-CD74 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.0) at 120°C for 2.5 min, UM800124) (1:800)




Immunohistochemical staining of paraffin-embedded Adenocarcinoma of Human ovary tissue using anti-CD74 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.0) at 120°C for 2.5 min, UM800124) (1:800)




Immunohistochemical staining of paraffin-embedded Human lymph node tissue within the normal limits using anti-CD74 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.0) at 120°C for 2.5 min, UM800124) (1:800)




Immunohistochemical staining of paraffin-embedded Human tonsil within the normal limits using anti-CD74 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.0) at 120°C for 2.5 min, UM800124) (1:800)




Immunohistochemical staining of paraffin-embedded NCI-H2228 cell pellets using anti-CD74 mouse monoclonal antibody (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.0) at 120°C for 2.5 min, UM800124) (1:800)




Immunohistochemical staining of paraffin-embedded MDA-MB-231 cell pellets using anti-CD74 mouse monoclonal antibody (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.0) at 120°C for 2.5 min, UM800124) (1:800)




Immunohistochemical staining of paraffin-embedded HL-60 cell pellets using anti-CD74 mouse monoclonal antibody (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.0) at 120°C for 2.5 min, UM800124) (1:800)




Immunohistochemical staining of paraffin-embedded LoVo cell pellets using anti-CD74 mouse monoclonal antibody (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.0) at 120°C for 2.5 min, UM800124) (1:800)




Immunohistochemical staining of paraffin-embedded mouse ascending colon tissue within the normal limits using anti-CD74 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA507339]) (1:400)



Immunohistochemical staining of paraffin-embedded mouse spleen tissue using anti-CD74 clone UMAB232 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, UM800124) (1:400).



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CD74 ([RC212964], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CD74 (1:500).



Western blot analysis of extracts (35ug) from 4 different cell lines and human spleen tissue by using anti-CD74 monoclonal antibody (1:500).