

Product datasheet for **UM800181**

Glutamine Synthetase (GLUL) Mouse Monoclonal Antibody [Clone ID: UMAB292]

Product data:

Product Type:	Primary Antibodies
Clone Name:	UMAB292
Applications:	IHC
Recommended Dilution:	IHC 1:15000
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human GS(GluI) (NP_001028228) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1.00mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Predicted Protein Size:	42.1 kDa
Gene Name:	glutamate-ammonia ligase
Database Link:	NP_001028228 Entrez Gene 2752 Human P15104



[View online »](#)

Background:


The protein encoded by this gene belongs to the glutamine synthetase family. It catalyzes the synthesis of glutamine from glutamate and ammonia in an ATP-dependent reaction. This protein plays a role in ammonia and glutamate detoxification, acid-base homeostasis, cell signaling, and cell proliferation. Glutamine is an abundant amino acid, and is important to the biosynthesis of several amino acids, pyrimidines, and purines. Mutations in this gene are associated with congenital glutamine deficiency, and overexpression of this gene was observed in some primary liver cancer samples. There are six pseudogenes of this gene found on chromosomes 2, 5, 9, 11, and 12. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Dec 2014]

Synonyms:


GLNS; GS; PIG43; PIG59

Protein Pathways:


Alanine, aspartate and glutamate metabolism, Arginine and proline metabolism, Metabolic pathways, Nitrogen metabolism

Product images:


IHC staining of FFPE human breast cancer tissue using anti-GS(Glul) mouse monoclonal antibody (UM800181) and Polink-2 HRP polymer detection kit ([D22-110]). Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 ([B04C-100]) at 120°C for 3 min. The brown stain indicates positive stain, blue is the counter stain.



IHC staining of FFPE human liver tissue with in the normal limits using anti-GS(Glul) mouse monoclonal antibody (UM800181) and Polink-2 HRP polymer detection kit ([D22-110]). Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 ([B04C-100]) at 120°C for 3 min. The brown stain indicates positive stain, blue is the counter stain.



IHC staining of FFPE human liver cancer tissue using anti-GS(Glul) mouse monoclonal antibody (UM800181) and Polink-2 HRP polymer detection kit (D22-110). Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 (B04C-100) at 120°C for 3 min. The brown stain indicates positive stain, blue is the counter stain.



IHC staining of FFPE human gastric tissue with in the normal limits using anti-GS(Glul) mouse monoclonal antibody (UM800181) and Polink-2 HRP polymer detection kit (D22-110). Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 (B04C-100) at 120°C for 3 min. The brown stain indicates positive stain, blue is the counter stain.