

Product datasheet for UM870050

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

DDX56 Mouse Monoclonal Antibody [Clone ID: UMAB153]

Product data:

Product Type: Primary Antibodies

Clone Name: UMAB153

Applications: 10k-ChIP, IF, IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:100~200

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 323-547 of human

DDX56 (NP_061955) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5~1.0 mg/ml (Lot Dependent)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: DEAD-box helicase 56

Database Link: NP 061955

Entrez Gene 54606 Human

Q9NY93




Background:

This gene encodes a member of the DEAD box protein family. DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA helicases. They are implicated in a number of cellular processes involving alteration of RNA secondary structure such as translation initiation, nuclear and mitochondrial splicing, and ribosome and spliceosome assembly. Based on their distribution patterns, some members of this family are believed to be involved in embryogenesis, spermatogenesis, and cellular growth and division. The protein encoded by this gene shows ATPase activity in the presence of polynucleotides and associates with nucleoplasmic 65S preribosomal particles. This gene may be involved in ribosome synthesis, most likely during assembly of the large 60S ribosomal subunit. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Mar 2012]


Synonyms: DDX21; DDX26; NOH61

Product images:




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human breast tissue using anti-DDX56 mouse monoclonal antibody. ([UM800050]; heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH9.0, 120°C for 3min)






Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human colon tissue using anti-DDX56 mouse monoclonal antibody. ([UM800050]; heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH9.0, 120°C for 3min)




Immunohistochemical staining of paraffinembedded Human Kidney tissue using anti-DDX56 mouse monoclonal antibody. ([UM800050]; heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH9.0, 120°C for 3min)




Immunohistochemical staining of paraffinembedded Carcinoma of Human kidney tissue using anti-DDX56 mouse monoclonal antibody. ([UM800050]; heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH9.0, 120°C for 3min)






Immunohistochemical staining of paraffinembedded Carcinoma of Human liver tissue using anti-DDX56 mouse monoclonal antibody. ([UM800050]; heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH9.0, 120°C for 3min)




Immunohistochemical staining of paraffinembedded Carcinoma of Human lung tissue using anti-DDX56mouse monoclonal antibody. ([UM800050]; heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH9.0, 120°C for 3min)




Immunohistochemical staining of paraffinembedded Human Ovary tissue using anti-DDX56 mouse monoclonal antibody. ([UM800050]; heatinduced epitope retrieval by 1mM EDTA in 10mM Tris, pH9.0, 120°C for 3min)






Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human ovary tissue using anti-DDX56 mouse monoclonal antibody. ([UM800050]; heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH9.0, 120°C for 3min)




Immunohistochemical staining of paraffinembedded Carcinoma of Human pancreas tissue using anti-DDX56 mouse monoclonal antibody. ([UM800050]; heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH9.0, 120°C for 3min)




Immunohistochemical staining of paraffinembedded Human thyroid tissue using anti-DDX56 mouse monoclonal antibody. ([UM800050]; heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH9.0, 120°C for 3min)






Immunohistochemical staining of paraffinembedded Carcinoma of Human thyroid tissue using anti-DDX56 mouse monoclonal antibody. ([UM800050]; heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH9.0, 120°C for 3min)




Immunohistochemical staining of paraffinembedded Human endometrium tissue using anti-DDX56 mouse monoclonal antibody. ([UM800050]; heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH9.0, 120°C for 3min)




Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human endometrium tissue using anti-DDX56 mouse monoclonal antibody. ([UM800050]; heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH9.0, 120°C for 3min)






Immunohistochemical staining of paraffinembedded Human prostate tissue using anti-DDX56 mouse monoclonal antibody. ([UM800050]; heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH9.0, 120°C for 3min)




Immunohistochemical staining of paraffinembedded Carcinoma of Human prostate tissue using anti-DDX56 mouse monoclonal antibody. ([UM800050]; heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH9.0, 120°C for 3min)




Immunohistochemical staining of paraffinembedded Human lymphoma tissue using anti-DDX56 mouse monoclonal antibody. ([UM800050]; heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH9.0, 120°C for 3min)






Western blot of human tissue lysates (15ug) from 10 different tissues (1: Testis, 2: Omentum, 3: Uterus, 4: Breast, 5: Brain, 6: Liver, 7: Ovary, 8: Thyroid, 9: Colon, 10: Spleen). Diluation: 1:500.




Western blot of mouse tissue lysates (20ug) from Spleen. Primary antibody diluation: 1:500. Secondary antibody dilution: Mouse TrueBlot® Ultra (1:1000).



Immunofluorescent staining of HeLa cells using anti-DDX56 mouse monoclonal antibody ([UM800050], green, 1:50). Actin filaments were labeled with Alexa Fluor® 594 Phalloidin (red), and nuclear with DAPI (blue). Scale bar, 20µm.





OriGene overexpression protein microarray chip was immunostained with UltraMAB anti-DDX56 mouse monoclonal antibody ([UM800050]). The positive reactive proteins are highlighted with two red arrows in the enlarged subarray. All the positive controls spotted in this subarray are also labeled for clarification.