

## Product datasheet for **UM870110**

### **GATA3 Mouse Monoclonal Antibody [Clone ID: UMAB218]**

#### **Product data:**


Product Type:	Primary Antibodies
Clone Name:	UMAB218
Applications:	10k-ChIP, IF, IHC, WB
Recommended Dilution:	IHC 1:50~100, IF 1:2000
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG2b
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 155-443 of human GATA3 (NP_002042) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.5~1.0 mg/ml (Lot Dependent)
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	47.9 kDa
Gene Name:	GATA binding protein 3
Database Link:	<a href="#">NP_001002295</a> <a href="#">Entrez Gene 14462 Mouse</a> <a href="#">Entrez Gene 85471 Rat</a> <a href="#">Entrez Gene 2625 Human</a> <a href="#">P23771</a>
Background:	This gene encodes a protein which belongs to the GATA family of transcription factors. The protein contains two GATA-type zinc fingers and is an important regulator of T-cell development and plays an important role in endothelial cell biology. Defects in this gene are the cause of hypoparathyroidism with sensorineural deafness and renal dysplasia. [provided by RefSeq, Nov 2009]


[View online »](#)


**Synonyms:** HDR; HDRS

**Protein Families:** Adult stem cells, ES Cell Differentiation/IPS, Stem cell relevant signaling - JAK/STAT signaling pathway, Transcription Factors


**Product images:**




Immunohistochemical staining of paraffin-embedded at margins of adenocarcinoma of human breast tissue using anti-GATA3 mouse monoclonal antibody. HIER pretreatment was done with 1mM EDTA in 10mM Tris buffer (pH8.0) at 120°C for 2.5 minutes. [UM800110] was diluted 1:50 and detection was done with HRP secondary and DAB chromogen. Strong nuclear stain is seen in the epithelial cells.




Immunohistochemical staining of paraffin-embedded human kidney tissue within the normal limits using anti-GATA3 mouse monoclonal antibody. HIER pretreatment was done with 1mM EDTA in 10mM Tris buffer (pH8.0) at 120°C for 2.5 minutes. [UM800110] was diluted 1:50, detection was done with HRP secondary and DAB chromogen. Weak nuclear stain is seen in the tubule epithelial cells of the kidney.




Immunohistochemical staining of paraffin-embedded carcinoma of human prostate tissue using anti-GATA3 mouse monoclonal antibody. HIER pretreatment was done with 1mM EDTA in 10mM Tris buffer (pH8.0) at 120°C for 2.5 minutes. [UM800110] was diluted 1:50 and detection was done with HRP secondary and DAB chromogen. Strong nuclear stain was seen in the tumor cells.




Immunohistochemical staining of paraffin-embedded human urothelial carcinoma using anti-GATA3 mouse monoclonal antibody. HIER pretreatment was done with 1mM EDTA in 10mM Tris buffer (pH8.0) at 120°C for 2.5 minutes. [UM800110] was diluted 1:50 and detection was done with HRP secondary and DAB chromogen. Strong nuclear stain is seen in the tumor cells.




Immunohistochemical staining of paraffin-embedded human bladder cancer using anti-GATA3 clone UMAB210 at 1:200 dilution of 1 mg/mL and detection with Polink2 Broad HRP DAB. [UM800110] requires heat-induced epitope retrieval with ACCEL (pH8.7) biocare pressure cooker at 110°C for 3minutes. The image shows nuclear staining in tumor cells.



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY GATA3 ([RC211904], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-GATA3 (1:500).




Western blot analysis of extracts (35ug) from 2 different cell by using anti-GATA3 monoclonal antibody (1:500).



Immunocytochemistry staining of MCF-7 cells using anti-GATA3 mouse monoclonal antibody ([UM800110]) (Left). The right is negative control. (1:2000)





OriGene overexpression protein microarray chip was immunostained with UltraMAB anti-GATA3 mouse monoclonal antibody ([UM800110]). The positive reactive proteins are highlighted with two red arrows in the enlarged subarray. All the positive controls spotted in this subarray are also labeled for clarification (1:100).