

## Product datasheet for **UM870111**


### **BOB1 (POU2AF1) Mouse Monoclonal Antibody [Clone ID: UMAB219]**

#### **Product data:**


Product Type:	Primary Antibodies
Clone Name:	UMAB219
Applications:	10k-ChIP, IHC
Recommended Dilution:	IHC 1:100~250
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Isotype:	IgG2b
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human POU2AF1(NP_006226) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	0.5~1.0 mg/ml (Lot Dependent)
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	27.3 kDa
Gene Name:	POU class 2 homeobox associating factor 1
Database Link:	<a href="#">NP_006226</a> <a href="#">Entrez Gene 18985 Mouse</a> <a href="#">Entrez Gene 690528 Rat</a> <a href="#">Entrez Gene 5450 Human</a> <a href="#">Q16633</a>
Background:	Transcriptional coactivator that specifically associates with either OCT1 or OCT2. It boosts the OCT1 mediated promoter activity and to a lesser extent, that of OCT2. It has no intrinsic DNA-binding activity. It recognizes the POU domains of OCT1 and OCT2. It is essential for the response of B-cells to antigens and required for the formation of germinal centers. [UniProtKB/Swiss-Prot Function]
Synonyms:	BOB1; OBF-1; OBF1; OCAB




[View online »](#)

**Product images:**


Immunohistochemical staining of paraffin-embedded human lymphoma tissue using anti-POU2AF1 mouse monoclonal antibody. HIER pretreatment was done with 1mM EDTA in 10mM Tris buffer (pH8.0) at 120°C for 2.5 minutes. [UM800111] was diluted 1:250 and detection was done with HRP secondary and DAB chromogen. Strong nuclear, cytoplasmic, and membranous staining tumor cells.




Immunohistochemical staining of paraffin-embedded Human tonsil within the normal limits using anti-POU2AF1 mouse monoclonal antibody. HIER pretreatment was done with 1mM EDTA in 10mM Tris buffer (pH8.0) at 120°C for 2.5 minutes. [UM800111] was diluted 1:250 and detection was done with HRP secondary and DAB chromogen. Strong nuclear, cytoplasmic, and membranous staining germinal cells of the tonsil.



Immunohistochemical staining of paraffin-embedded tonsil using mouse anti-POU2AF1 clone UMAB219 ([UM800111]) at 1:200 with Polink2 Broad HRP DAB detection kit; pretreatment of tissue prior to stain with heat-induced epitope retrieval with EDTA pH 8.0 buffer using pressure chamber for 3 minutes at 110C is required for optimal staining. Cells exhibit cytoplasmic and membranous staining.



Immunohistochemical staining of paraffin-embedded normal spleen using mouse anti-POU2AF1 clone UMAB219 ([UM800111]) at 1:200 with Polink2 Broad HRP DAB detection kit; pretreatment of tissue prior to stain with heat-induced epitope retrieval with EDTA pH 8.0 buffer using pressure chamber for 3 minutes at 110C is required for optimal staining. Cells exhibit cytoplasmic and membranous staining.



OriGene overexpression protein microarray chip was immunostained with UltraMAB anti-POU2AF1 mouse monoclonal antibody ([UM800111]). The positive reactive proteins are highlighted with two red arrows in the enlarged subarray. All the positive controls spotted in this subarray are also labeled for clarification (1:100).