

Product datasheet for UM870155

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CDX2 Mouse Monoclonal Antibody [Clone ID: UMAB265]

Product data:

Product Type: Primary Antibodies

Clone Name: UMAB265

Applications: IHC

Recommended Dilution: IHC 1:300

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human CDX2 (NP_001256) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5~1.0 mg/ml (Lot Dependent)

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 33.3 kDa

Gene Name: caudal type homeobox 2

Database Link: NP 001256

Entrez Gene 12591 MouseEntrez Gene 66019 RatEntrez Gene 1045 Human

Q99626

Background: Involved in the transcriptional regulation of multiple genes expressed in the intestinal

epithelium. Important in broad range of functions from early differentiation to maintenance of the intestinal epithelial lining of both the small and large intestine. [UniProtKB/Swiss-Prot

Function]

Synonyms: CDX-3; CDX2/AS; CDX3







Protein Families:

Adult stem cells, Druggable Genome, Embryonic stem cells, ES Cell Differentiation/IPS, Transcription Factors

Product images:



Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human colon tissue using anti-CDX2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.0) at 120°C for 2.5 min, [UM800155]) (1:300)



Immunohistochemical staining of paraffinembedded Human appendix tissue within the normal limits using anti-CDX2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.0) at 120°C for 2.5 min, [UM800155]) (1:300)